

Mouse Monoclonal Antibody to P16 (Mouse and Human)

Order Information				
Catalog#	20129			
Size/Concentration	50ul	100µl		
Price(¥)	1280	2180		

Description

The progression of cells through the cell cycle is regulated by a family of protein kinases known as cyclin-dependent kinases (Cdks). The sequential activation of individual members of this family and their consequent phosphorylation of critical substrates promotes orderly progression through the cell cycle. The cyclins function as differentially expressed positive regulators of Cdks. Negative regulators of the cycle include the p53-inducible 21 kDa WAF1/Cip1 protein designated p21, Kip1 p27 and p16. The complexes formed by Cdk4 and the D-type cyclins have been strongly implicated in the control of cell proliferation during the G1 phase. It has recently been shown that p16 binds to Cdk4 and inhibits the catalytic activity of the Cdk4/cyclin D complex. Moreover, the gene encoding p16 exhibits a high frequency of homozygous deletions and point mutations in established human tumor cell lines. Call 1-510-860-4615 +86-19375157864 Email Info@ProMab.com Web www.ProMab.com www.ProMab.cn

Protocal

WB - www.promab.com/protocol/wb.html IHC - www.promab.com/protocol/ihc.html ICC - www.promab.com/protocol/icc.html HCM - www.promab.com/protocol/hcm.html **Antigen Sequence** is available upon request.


Specification			
Entrez GenelD	1029		
Aliases	P16		
Clone#	2D9A12		
Host/Isotype	Mouse IgG2b		
Storage	4°C; -20°C for long term storage. Avoid freeze /thaw cycles.		
Species Reactivity	Human,Rat		
Immunogen	Purified recombinant fragment of P16 express ed in E. Coli.		
Formulation	Ascitic fluid containing 0.03% sodium azide.		

Application		
ELISA	1/10000	
IHC	1/200 - 1/1000	

References


- 1. Hunter, T. 1993. Cell 75: 839-841.
- 2. Sherr, C.J. 1993. Cell 73: 1059-1065.
- 3. El-Deiry, W.S., et al. 1993. Cell 75: 817-825.

Products and Services





Mouse Monoclonal Antibody to P16 (Mouse and Human)



Immunohistochemical analysis of paraffin-embedded rat liver tissue (A), human brain tissue (B) and brain tumor (C), showing nuclear localization using P16 mouse mAb with DAB staining.

#20129