

Mouse Monoclonal Antibody to ACHE

Order Information

Catalog#	32394	
Size/Concentration	50ul	100ul
Price(¥)	1280	2180

Description

Acetylcholinesterase hydrolyzes the neurotransmitter, acetylcholine at neuromuscular junctions and brain cholinergic synapses, and thus terminates signal transmission. It is also found on the red blood cell membranes, where it constitutes the Yt blood group antigen. Acetylcholinesterase exists in multiple molecular forms which possess similar catalytic properties, but differ in their oligomeric assembly and mode of cell attachment to the cell surface. It is encoded by the single ACHE gene, and the structural diversity in the gene products arises from alternative mRNA splicing, and post-translational associations of catalytic and structural subunits. The major form of acetylcholinesterase found in brain, muscle and other tissues is the hydrophilic species, which forms disulfide-linked oligomers with collagenous, or lipid-containing structural subunits. The other, alternatively spliced form, expressed primarily in the erythroid tissues, differs at the C-terminal end, and contains a cleavable hydrophobic peptide with a GPI-anchor site. It associates with the membranes through the phosphoinositide (PI) moieties added post-translationally. AChE activity may constitute a sensitive biomarker of RBC ageing in vivo, and thus, may be of aid in understanding the effects of transfusion.

Specification

Entrez GeneID	43
Aliases	YT; ACEE; ARACHE; N-ACHE
Clone#	3D5E8
MW	67.8kDa
Host/Isotype	Mouse IgG1
Storage	4°C; -20°C for long term storage. Avoid freeze/thaw cycles.
Species Reactivity	Human, Mouse, Monkey, Rat
Immunogen	Purified recombinant fragment of human AChE expressed in E. Coli.
Formulation	Purified antibody in PBS with 0.05% sodium azide

Application

WB	1/500 - 1/2000
FCM	1/200 - 1/400
IHC	1/200 - 1/1000

References

1. Molecules. 2020 Oct 4;25(19):4545.
2. J Cell Biochem. 2021 Aug;122(8):787-800.






Call 1-510-860-4615
+86-19375157864
Email Info@ProMab.com
Web www.ProMab.com
www.ProMab.cn

Protocol

WB - www.promab.com/protocol/wb.html
IHC - www.promab.com/protocol/ihc.html
ICC - www.promab.com/protocol/icc.html
HCM - www.promab.com/protocol/hcm.html

Antigen Sequence is available upon request.

Products and Services


-  Mouse Monoclonal Antibody
-  Rat Monoclonal Antibody
-  Human Antibody
-  Hybridoma Sequencing
-  Polyclonal Antibody

For Research Only


Application Key: WB - Western Blot | IHC - Immunohistochemistry | ICC - Immunocytochemistry | FCM - Flow Cytometry | ELISA - Enzyme-linked Immunosorbent Assay | IP - Immunoprecipitation

#32394


Mouse Monoclonal Antibody to ACHE




Western blot analysis using ACHE mouse mAb against PC-12 (1), HeLa (2), mouse brai (3), NIH/3T3 (4), COS7 (5), Jurkat (6) and Raji (7) cell lysate.



Flow cytometric analysis of NIH/3T3 cells using ACHE mouse mAb (green) and negative control (red).



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissues using ACHE mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded human rectum cancer tissues using ACHE mouse mAb with DAB staining.

For Research Only

Application Key: **WB** - Western Blot | **IHC** - Immunohistochemistry | **ICC** - Immunocytochemistry | **FCM** - Flow Cytometry | **ELISA** - Enzyme-linked Immunosorbent Assay | **IP** - Immunoprecipitation

#32394